

Part I: obs4MIPs (P. Gleckler) Part II: input4MIPs (P. Durack)

ESGF F2F 2017, Sheraton Fisherman's Wharf, San Francisco

Tuesday 5th December 2017

PROGRAM FOR CLIMATE MODEL DIAGNOSIS AND INTERCOMPARISON (PCMDI) pcmdi.llnl.gov/home







An Update on obs4MIPs from an ESGF perspective: progress plans and challenges

WDAC Observations for Model Evaluation Task Team

Peter Gleckler, co-chair (PCMDI) and Duane Waliser, co-chair (JPL/NASA),
Mike Bosilovich (GSFC/NASA), Helene Chepfer (IPSL), Veronika Erying (DLR), Robert Ferraro (JPL/NASA),
Pierre-Phillipe Mathieu (ESA), Jerry Potter (GSFC), Roger Saunders (UKMO),
Jörg Schulz (EUMETSAT), Karl Taylor (PCMDI), Jean-Noël Thépaut (ECMWF)

Additional regular contributors:

Otis Brown (WDAC), Michel Rixen (WCRP), Sophie Cloché (IPSL), Tsengdar Lee (NASA) and Renu Joseph (DOE),
Luca Cinquini (JPL) – CoG technical support
Denis Nadeau (PCMDI) – CMOR3 development, Paul Durack (PCMDI) - obs4MIPs controlled vocabulary,
Jim Biard (NCEI) and Matthias Tuma (WCRP)


... and others


This work was supported by the U.S. Department of Energy's (DOE's) Office of Science (Biological and Environmental Research) through its Regional and Global Climate Modeling Program and was performed at Lawrence Livermore National Laboratory as a contribution to the U.S. Department of Energy, Office of Science, Climate and Environmental Sciences Division, Regional and Global Climate Modeling Program under Contract DE-AC52-07NA27344.

obs4MIPs

https://www.earthsystemcog.org/projects/obs4mips/

- A Project for identifying, documenting and disseminating observations for climate model evaluation.
- Data sets accessible on the ESGF alongside CMIP model output, adhering to the same data conventions
- Guided by the WCRP Data Advisory Council obs4MIPS Task Team



















Progress since ESGF F2F #6

(the not so good news)

New entries to obs4MIPs essentially stalled, awaiting...

CMIP6 global attributes, DRS, filenames, directory structure and CVs
 Updating obs4MIPs data specifications to match CMIP6
 Management of obs4MIPs CMOR tables and CVs on github
 CMOR refinements
 process documentation, beta testing
 ready for data providers

Progress since ESGF F2F #6

(more recently... the good news!)

- CMIP6 global attributes, DRS, filenames, directory structure and CVs \checkmark
 - obs4MIPs data specifications* (ODS 2.1) √
- Management of obs4MIPs CVs on github √
- Additional CMOR enhancements to deal with gridded observations ✓
 (now being used for new NOAA and ESA contributions)
- Establishment of two new features √
 - Dataset "suitability indicators" √
 - Introduction of "supplemental information" √

^{*} Includes placeholders for ES-DOC and DKRZ publication service

obs4MIPs Dataset Suitability & Maturity Indicators

Dataset


Comparison

Technical

Requirements		Suitability and Maturity			Complexity
Meets obs4MIPs data technical requirements	Includes obs4MIPs technical note information	Closeness or robustness of measurement to observed reference quantity	Maturity with respect to climate model evaluation	Provision for robust uncertainty information	Complexity of Model Observation Comparison
Data suitably processed with CMOR and/or consistent with obs4MIPs standards	Complete technical note information provided	Measurement approach provides a very close relationship to observation quantity	Multiple peer-reviewed examples of application to climate model evaluation	Uncertainty information provided per retrieval/grid point	Comparison can be made directly with CMIP model output variable
Largely complete with minor metadata inconsistencies	Technical note information incomplete and/or could be improved	Measurement approach requires complex and/or non-linear retrieval methods and/or subjective inferences/definitions	One peer-reviewed example of application to climate or component model evaluation.	General uncertainty information given relative to the methodology and dataset as a whole - backed by actual field/in- situ validation exercises	Comparison requires some simple post processing of CMIP output variable(s) (e.g. vertical integral or ratio of two variables)
Non-compliant. Should be removed from database!	Technical note not provided	Measurement approach requires significant use/influence from complex or weakly constrained model and/or has significant ambiguity in definition(s)	No peer-reviewed xamples of application to model evlauation	No uncertainty information provided	Comparison requires complex processing of CMIP output (e.g. "simulator", budget calculation)

Illustration Of Dataset Suitability And Maturity Indicators


Based on typical obs4MIPs dataset search



Linked as a single entity to explanatory page of color codes

Illustration Of Dataset Suitability And Maturity Indicators

Based on typical obs4MIPs dataset search



Access to SUPPLEMENTAL DATA (in zip file; instructions available)

Example: obs4MIPs "registered content": (Data providers enter as github issue)

```
"NOAA-NCEI-ERSST-4-0":{
    "institution_id":"NOAA-NCEI",
    "region":[
        "global ocean"
    "release year": "2015",
    "source_description": "Extended Reconstructed Sea Surface Temperatures",
    "source_id": "NOAA-NCEI-ERSST-4-0",
    "source label": "NOAA-NCEI-ERSST",
    "source name": "NOAA NCEI ERSST",
    "source type": "gridded insitu",
    "source variables":[
        "tos"
    ],
    "source version number": "4.0"
},
```

Questions...

Q: What are the key things that are difficult to do today and are impeding scientific progress or productivity?

A: Preparing and publishing data

Q: What are key development effort that you see are needed for the future success of your projects?

A: Streamlining the process of contributing data

Q: What is the estimated size of your distributed archive?

A: ~Terabytes

Q: What are the administrative/sponsor requirements that arise from your project?

A: Usage metrics, persistent and digital object identifiers, dataset version control

Q: What are your expected strategic roadmaps for the ESGF's short, mid and long term development efforts?

A: 1-3 years: Get caught up, including updating existing datasets, with new data specifications (ODS 2.1)

3-5 years: Get a majority of gridded data used for model evaluation research in obs4MIPs

5-10 years: It is possible, with sufficient progress, that obs4MIPs or a similar effort will revolutionize how data are coordinated across agencies, borders, and classes of data

obs4MIPs: Current ESGF strategic needs

- Smooth transition to new facets/datasets;
 backwards compatibility with existing data
- Remove or filter unwanted data (by whom?)
- Planning for in-situ data



input4MIPs: Boundary Conditions and Forcing Datasets for CMIP6

DOE ESGF F2F 2017, Sheraton Fisherman's Wharf, San Francisco

Paul J. Durack, Karl E. Taylor, Sasha Ames, Tony Hoang, Martina Stockhause and many others.. Tuesday 5th December 2017

PROGRAM FOR CLIMATE MODEL DIAGNOSIS AND INTERCOMPARISON (PCMDI) pcmdi.llnl.gov/home









Section 0: What is input4MIPs?

Observationally-derived input datasets for Model Intercomparison Projects (MIPs)





- Hosts all required CMIP (DECK) and satellite MIP experiments forcing datasets
- Currently ~1200 files, ~0.5TB, ~15 contributing institutions
- User base is small ~100 users, ~30+ modeling groups






https://esaf-node.linl.gov

ESGE P2P Version v2 5 13 master


NOAA | NASA | NSF | DoE Office of Science | IS-ENES

Moving target – using "live" google doc to keep track of changing datasets






- Search facets have attempted to conform as closely to CMIP6 as possible
- Learning-by-doing publication quirks will be fixed in a Jan '18 republication





- Search facets have attempted to conform :
- Learning-by-doing publication quirks will be



Variable

Target MIP

+



Section 1: Problem data

Expanding ESGF support for more data formats



- Large single files vs temporal chunking (1 x 16GB vs 8 x 2GB time chunks)
- ~16GB (UMD) user downloads in 3rd world countries

New! LUH2 Historical datasets now available

LUH2 v2h Release (10/14/16): The updated release of the historical land-use forcing dataset (LUH2 v2h) covers the period 850-2015 and corrects all known issues and notices identified with the previous version (LUH2 v1.0h). This dataset replaces the previously released dataset (LUH2 v1.0h). This product is the result of a series of prototypes released previously, uses the established data format, and will connect smoothly to gridded products for the future.

- Historic Data (850 2015 AD)
 - states.nc (5.8 CP)
 - o transitions. (16 GB)
 - o management.ne (1.4 GB)
- Supporting Files
 - staticData quarterdeg.nc (1 MB)
- Data Documentation
 - LUH2 v2h README
- Will large files be an issue for users behind firewalls?
- What is the experience with download restarts with wget (what about rsync?)



Non-CMIP data: Multiple variables per file (UoM) – rewrite if possible

```
[durack1@oceanonly CMIP6]$ ncdump -h input4MIPs/UoM/GHGConc/CMIP/yr/atmos/UoM-CMIP-1-1-0/GHGCon
c/gr3-GMNHSH/v20160701/mole_fraction_of_carbon_dioxide_in_air_input4MIPs_GHGConcentrations_CMIP
_UoM-CMIP-1-1-0_gr3-GMNHSH_0000-2014.nc
netcdf mole_fraction_of_carbon_dioxide_in_air_input4MIPs_GHGConcentrations_CMIP_UOM-CMIP-1-1-0_
gr3-GMNHSH_0000-2014 {
dimensions:
       lat = 1 :
       bnds = 2;
       time = 2015;
variables:
       double lat(lat);
                lat:units = "degrees_north" ;
                lat:long name = "latitude" ;
                lat:standard name = "latitude" ;
                lat:axis = "Y" ;
                lat:bounds = "lat_bnds" ;
       double lat_bnds(lat, bnds);
       double time(time) ;
                time:units = "days since 1850-01-01 00:00:00";
               time:calendar = "365_day";
               time:long_name = "time";
               time:standard_name = "time";
               time:bounds = "time_bnds" ;
       double time_bnds(time, bnds);
       float carbon_dioxide_GM(time) ;
               carbon_dioxide_GM:long_name = "Global Mean Mole Fraction of CO2" ;
               carbon_dioxide_GM:original_name = "CO2_GM" ;
               carbon dioxide GM:standard_name = "mole_fraction_of_carbon_dioxide_in_air";
               carbon_dioxide_GM:units = "1.e-6" ;
               carbon_dioxide_GM:cell_methods = "time: mean area: mean" ;
               carbon_dioxide_GM:lat = "0.0";
               carbon_dioxide_GM:lat_bnds = "-90.0, 90.0";
       float carbon dioxide NH(time);
               carbon_dioxide_NH:long_name = "Northern Hemisphere Mean Mole Fraction of CO2";
               carbon_dioxide_NH:original_name = "CO2_NH" ;
               carbon_dioxide_NH:standard_name = "mole_fraction_of_carbon_dioxide_in_air" ;
               carbon dioxide NH:units = "1.e-6";
               carbon_dioxide_NH:cell_methods = "time: mean area: mean" ;
               carbon_dioxide_NH:lat = "30.0";
               carbon_dioxide_NH:lat_bnds = "0.0, 90.0";
       float carbon dioxide SH(time) :
               carbon_dioxide_SH:long_name = "Southern Hemisphere Mean Mole Fraction of CO2";
               carbon_dioxide_SH:original_name = "CO2_SH" ;
               carbon_dioxide_SH:standard_name = "mole_fraction_of_carbon_dioxide_in_air" ;
               carbon_dioxide_SH:units = "1.e-6"
                carbon_dioxide_SH:cell_methods = "time: mean area: mean" ;
               carbon_dioxide_SH:lat = "-30.0";
               carbon_dioxide_SH:lat_bnds = "-90.0, 0.0";
// global attributes:
                :title = "UoM-CMIP-1-1-0: historical GHG concentrations: global and hemispheric
 means of CO2 prepared for input4MIPs";
               :institution_id = "UoM";
               :dataset_category = "GHGConcentrations";
                :dataset_version_number = "1.1.0";
```



:title = "UoM-CMIP-1-1-0: historical GHG concentrations: global and hemispheric

// global attributes:

means of CO2 prepared for input4MIPs";

:institution_id = "UoM";


:dataset_category = "GHGConcentrations" ;
:dataset_version_number = "1.1.0" ;

Non-CMIP data: Multiple variables per file (UoM) – rewrite if possible

```
[durack1@oceanonly CMIP6]$ ncdump -h input4MIPs/UoM/GHGConc/CMIP/yr/atmos/UoM-CMIP-1-1-0/GHGCon
c/gr3-GMNHSH/v20160701/mole_fraction_of_carbon_dioxide_in_air_input4MIPs_GHGConcentrations_CMIP
_UoM-CMIP-1-1-0_gr3-GMNHSH_0000-2014.nc
netcdf mole_fraction_of_carbon_dioxide_in_air_input4MIPs_GHGConcentrations_CMIP_UoM-CMIP-1-1-0_
                                                                                                             Rewritten file...
gr3-GMNHSH_0000-2014 {
dimensions:
       lat = 1 :
                                                                          [durack1@oceanonly CMIP6]$ ncdump -h input4MIPs/UoM/GHGConcentrations/CMIP/yr/atmos/UoM-CMIP-1-
       bnds = 2;
                                                                          2-0/mole fraction of carbon dioxide in air/qr1-GMNHSH/v20160830/mole fraction of carbon dioxide
       time = 2015:
                                                                          in air input4MIPs GHGConcentrations CMIP UoM-CMIP-1-2-0 gr1-GMNHSH 0000-2014.nc
variables:
                                                                          netcdf mole_fraction_of_carbon_dioxide_in_air_input4MIPs_GHGConcentrations_CMIP_UoM-CMIP-1-2-0_
       double lat(lat);
              lat:units = "degrees_north";
                                                                          gr1-GMNHSH 0000-2014 {
               lat:long name = "latitude" ;
                                                                          dimensions:
               lat:standard name = "latitude" ;
                                                                                   time = UNLIMITED ; // (2015 currently)
              lat:axis = "Y" ;
                                                                                  bound = 2;
               lat:bounds = "lat_bnds" ;
                                                                                   sector = 3;
       double lat_bnds(lat, bnds);
                                                                          variables:
       double time(time) ;
               time:units = "days since 1850-01-01 \ 00:00:00";
                                                                                   float time(time);
              time:calendar = "365_day";
                                                                                           time:bounds = "time bnds" :
              time:long_name = "time";
                                                                                           time:long_name = "time";
              time:standard_name = "time" ;
                                                                                           time:standard_name = "time";
                                                                                           time:units = "days since 0-1-1";
              time:bounds = "time_bnds" ;
                                                                                           time:calendar = "gregorian" ;
       double time_bnds(time, bnds);
                                                                                           time:axis = "T" :
       float carbon_dioxide_GM(time) ;
                                                                                   double time_bnds(time, bound) ;
              carbon_dioxide_GM:long_name = "Global Mean Mole Fraction of CO
                                                                                   int sector(sector);
              carbon dioxide GM:original name = "CO2 GM";
              carbon dioxide GM:standard_name = "mole_fraction_of_carbon_dio"
                                                                                           sector:bounds = "sector bnds";
              carbon_dioxide_GM:units = "1.e-6" ;
                                                                                           sector:lat_bnds = "0: -90.0, 90.0; 1: 0.0, 90.0; 2: -90.0, 0.0";
              carbon_dioxide_GM:cell_methods = "time: mean area:
                                                                                           sector:long_name = "sector";
              carbon_dioxide_GM:lat = "0.0";
                                                                                           sector:ids = "0: Global; 1: Northern Hemisphere; 2: Southern Hemisphere";
              carbon_dioxide_GM:lat_bnds = "-90.0, 90.0";
                                                                                           sector:original names = "0: CO2 GM; 1: CO2 NH; 2: CO2 SH";
       float carbon dioxide NH(time);
                                                                                  double sector_bnds(sector, bound) ;
              carbon dioxide NH:long name = "Northern Hemisphere Mean Mole F
                                                                                   float mole_fraction_of_carbon_dioxide_in_air(time, sector);
              carbon_dioxide_NH:original_name = "CO2_NH" ;
                                                                                           mole_fraction_of_carbon_dioxide_in_air:_FillValue = 1.e+20f ;
              carbon_dioxide_NH:standard_name = "mole_fraction_of_carbon_dio
                                                                                           mole_fraction_of_carbon_dioxide_in_air:missing_value = 1.e+20f ;
              carbon dioxide NH:units = "1.e-6";
              carbon_dioxide_NH:cell_methods = "time: mean area: mean";
                                                                                           mole_fraction_of_carbon_dioxide_in_air:long_name = "mole";
              carbon_dioxide_NH:lat = "30.0";
                                                                                           mole_fraction_of_carbon_dioxide_in_air:cell_methods = "time: mean area: mean" ;
              carbon_dioxide_NH:lat_bnds = "0.0, 90.0";
                                                                                           mole fraction of carbon dioxide in air:units = "1.e-6";
       float carbon dioxide SH(time) :
              carbon_dioxide_SH:long_name = "Southern Hemisphere Mean Mole"
                                                                           // global attributes:
              carbon_dioxide_SH:original_name = "CO2_SH";
                                                                                           :Conventions = "CF-1.6" :
              carbon_dioxide_SH:standard_name = "mole_fraction_of_carbon_diox
              carbon_dioxide_SH:units = "1.e-6" ;
                                                                                           :comment = "Data provided are global and hemispheric area-weighted means. Zonal
              carbon_dioxide_SH:cell_methods = "time: mean area: mean";
                                                                           means for 15-degree lat bands or 0.5-degree lat bands are available in gn-15x360 or gr-0p5x360
              carbon_dioxide_SH:lat = "-30.0";
                                                                           files respectively";
              carbon dioxide SH:lat bnds = "-90.0, 0.0";
                                                                                           :variable id = "mole fraction of carbon dioxide in air" :
```


PCMDI

Non-netcdf – tar gzipped, code snippets, text files (MPI)





Non-netcdf – tar gzipped, code snippets, text files (MPI-M)



ESGF P2P Version v2.5.13 master



NOAA I NASA I NSF I DoE Office of Science I IS-ENES

Privacy & Legal Notic

DoE Office of Science I IS-ENES I NASA I NOAA I NCI I NSF

Section 2:


Deprecating datasets

How to gracefully "hide" obsolete datasets, but leave them available for trace-ability



Deprecating datasets (gracefully):


- Useful for datasets that are deprecated (rather than unpublishing)
- PCMDI AMIP dataset is updated every 6 months 1.1.0 vs 1.1.1 vs 1.1.2 vs 1.1.3..





Deprecating datasets (gracefully):

- Useful for datasets that are deprecated (rather than unpublishing)
- PCMDI AMIP dataset is updated every 6 months 1.1.0 vs 1.1.1 vs 1.1.2 vs 1.1.3..





Section 3:

DOI minting and DKRZ citation service

Making contributed data findable and citable



DKRZ Citation Service active:

Useful for dataset discovery and citation - brings recognition to data providers

PCMDI AMIP dataset is updated every 6 months — 1.1.0 vs 1.1.1 vs 1.1.2 vs

1.1.3..



http://doi.org/ 10.22033/ESGF/ input4MIPs.1161



Section 4:


Integrated with the ES-DOC model documentation system

Making input4MIPs versioned data correctly citable by modeling groups




Keeping track of data versions:

- Contributed datasets have been growing
- Multiple versions since December 2016, with 3 CMIP panel "official" releases and a multitude of updates in between official releases



Keeping track of data versions:

- Contributed datasets have been growing
- Multiple versions since December 2016, with 3 CMIP panel "official" releases and a multitude of updates (~27 versions) between official releases
- Attempt to manage version info using json dictionaries for use by ES-DOCs





Section 5:

Ensuring consistent publication metadata across the project



If data doesn't follow identical metadata conventions:

Maintaining consistent publication entries is a

challenge

```
b UofMD vr UofMD-landState-MAGPIE-ssp585-2-1-f_multiple-transitions.json (~/Desktop) - VIM
    "input4MIPs-RitvikSahajpal_LandUseForcing":{
        "Conventions": "CF-1.6",
        "activity_id":"input4MIPs",
        "contact": gchurtt@umd.edu, lchini@umd.edu, steve.frolking@unh.edu, ritvik@umd.edu",
        "creation_date":"2017-09-16T23:15:45Z",
        "dataset_category":"landState",
        "dataset version_number": "2.1f",
        "file":"CMIP6/input4MIPs/UofMD/landState/ScenarioMIP/yr/land/UofMD-landState-MAGPIE-ss
p585-2-1-f/multiple-transitions/gn/v20171005/multiple-transitions_input4MIPs_landState_Scenari
oMIP UofMD-MAGPIE-ssp585-2-1-f gn 2015-2100.nc",
        "frequency": "yr",
        "further_info_url": "http://luh.umd.edu",
        "grid label": "gn",
        "institution": "University of Maryland (UofMD), College Park, MD 20742, USA",
        "institution_id":"UofMD",
        "mip era":"CMIP6",
        "nominal_resolution":"25 km",
        "project": "input4MIPs",
        "realm":"land",
        "source": "LUH2 v2.1f: Land-Use Harmonization Data Set",
        "source id": "UofMD-landState-MAGPIE-ssp585-2-1-f",
        "target mip":[
            "ScenarioMIP",
            "AerChemMIP",
            "C4MIP",
            "DCPP",
            "LS3MIP",
        "timestamp": "2017-10-05T18:29:23Z",
        "title": "UofMD LUH2f dataset prepared for input4MIPs",
        "tracking_id":"hdl:21.14100/c7bb2ffe-a32a-4fb3-b742-196f3bc89661",
        "variable_id": "multiple-transitions",
        "version":"v20171005"
```

Section 6:

Replication with SYNDA

Multiple variable files do not conform to CMIP-standards and..



SYNDA replication:

- SYNDA has hit problems downloading input4MIPs data with more than a single variable per file
- Contributed data passed the CF-check, but isn't CMIP "format" single variable per file
- input4MIPs data isn't derived from models, the DRS structure differs from the CMIP data
- Many of the "obs" products contributed have interacting variables – to use the data you need more than a single variable in memory
- *esg.ini configuration changes seem to have solved the issue



Thank you

Peter J. Gleckler t +1 925 422 7631 e gleckler1@llnl.gov w pcmdi.llnl.gov Paul J. Durackt +1 925 422 5208e pauldurack@llnl.govw pcmdi.llnl.gov

